



Saskatchewan
Ministry of
Agriculture

Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 16, July 19, 2012

For the Period July 10 to 16, 2012

Saskatchewan livestock producers have 65 per cent of the 2012 hay crop cut and 42 per cent baled or put into silage, according to Saskatchewan Agriculture's weekly Crop Report. Ninety-one per cent of the hay crop is rated as good to excellent in quality.

The estimated average hay yields on dry land are reported as 1.4 tons per acre (alfalfa and wild hay), 1.6 tons per acre (alfalfa/brome and other tame hay) and 1.9 tons per acre (greenfeed). On irrigated land, the estimated average hay yields are 2.1 tons per acre (alfalfa), 2.3 tons per acre (alfalfa/brome), 2.4 tons per acre (other tame hay) and 2.6 tons per acre (greenfeed).

Warm and dry weather this past week has allowed for crops to advance quickly and most of them are in good condition. High temperatures and humidity have resulted in thunderstorms in some areas that produced varying amounts of precipitation. Disease, insects and wind are causing the majority of crop damage.

Cropland top soil moisture is rated as 18 per cent surplus, 74 per cent adequate and eight per cent short. Hay land and pasture topsoil moisture is rated as 11 per cent surplus, 75 per cent adequate, 13 per cent short and one per cent very short.

Farmers are busy haying and controlling diseases and insects.

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

Crop development is quickly advancing due to warm temperatures. Most of the area experienced high humidity that resulted in thunderstorms and varying amounts of rainfall this past week. Heavy rain and strong winds have damaged many cereal crops and some fields are severely lodged. The Frobisher area reported 26 mm of rainfall, the Moosomin area 20 mm, the Rocanville area 12 mm, the Weyburn area 15 mm, the Marquis area 28 mm and the Ceylon area 8 mm. The Tantallon area has recorded the greatest amount of precipitation since April 1 (476 mm).

For further information, contact: Grant McLean, PAG,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

One year ago

Fifty-nine per cent of the hay crop had been cut with 34 per cent baled or put into silage.

Eighty-seven per cent of the hay crop was rated as good to excellent in quality.

Follow the 2012 Crop Report
on Twitter [@SKGovAg](https://twitter.com/SKGovAg)

Estimated Provincial Hay Yields (tons/acre) - July 16, 2012

	Dry land	Irrigated Land
Alfalfa	1.4	2.1
Brome/Alfalfa	1.6	2.3
Other Tame Hay	1.6	2.4
Wild Hay	1.4	N/A
Greenfeed	1.9	2.6



Agriculture and Agri-Food Canada Agriculture et Agroalimentaire Canada

SCIC

SASKATCHEWAN CROP
INSURANCE CORPORATION

Topsoil moisture conditions on cropland are rated as 17 per cent surplus, 71 per cent adequate and 12 per cent short. Hay land and pasture moisture is rated as eight per cent surplus, 74 per cent adequate, 17 per cent short and one per cent very short. Crop districts 1B and 2B are reporting 20 per cent and 30 per cent of acres, respectively, to have surplus topsoil moisture. Crop districts 2A and 3ASE are reporting 38 per cent and 43 per cent of respective acres to be short of topsoil moisture.

Estimated Southeast Hay Yields (tons/acre) - July 16, 2012		
	Dry land	Irrigated Land
Alfalfa	1.1	N/A
Brome/Alfalfa	1.5	N/A
Other Tame Hay	1.4	N/A
Wild Hay	1.3	N/A
Greenfeed	1.5	N/A

Livestock producers have 75 per cent of the hay crop cut and 56 per cent has been baled or put into silage. Hay quality is rated as 13 per cent excellent, 83 per cent good and four per cent fair. Alfalfa weevils have damaged many pure stands and hay yield has been reduced.

The majority of crop damage is due to wind, disease, flooding and drought. Many crops in the area experienced heat stress, and some cooler temperatures and rainfall would be appreciated to help the crops recover. Some producers are applying pesticides for leaf disease, fusarium head blight and wheat midge. Producers are busy haying and controlling disease and insects in crops.

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

Precipitation in the area ranged from nil to 49 mm, with the majority of producers receiving very little rainfall. The Lisieux received 15 mm of rain, the Fife Lake area 3 mm, the Mortlach area 49 mm, the Chaplin area 8 mm, the Vanguard area 7 mm, the Swift Current area 5 mm

Estimated Southwest Hay Yields (tons/acre) - July 16, 2012		
	Dry land	Irrigated Land
Alfalfa	1.3	1.7
Brome/Alfalfa	1.4	1.8
Other Tame Hay	2.2	2.2
Wild Hay	1.7	N/A
Greenfeed	1.5	2.0

and the Consul area 20 mm. Cropland topsoil moisture is rated as 72 per cent adequate, 27 per cent short and one per cent short. Hay land and pasture topsoil moisture is rated as 69 per cent adequate, 27 short and four per cent very short. Many crop districts are reporting a large number of cropland, hay land and pasture acres to be short of moisture.

Livestock producers have 75 per cent of the hay crop cut and 49 per cent has been baled or put into silage. Hay crop quality is rated as 18 per cent excellent, 66 per cent good, 13 per cent fair and three per cent poor.

High temperatures and humidity have stressed many crops, and additional rainfall is needed to help them through the rest of summer. Drought, wind, insects and disease have caused the majority of crop damage. Crop development has advanced quickly and there are indications that winter cereals and some pulses will be ready to be harvested in the coming weeks.

Farmers are busy haying, controlling disease in crops and getting harvest equipment ready.

East-Central Saskatchewan (Crop Districts 5 and 6a)

Large amounts of rainfall were received in most of the area this past week. The Roblin area recorded 51 mm of rain, the Meacham area 43 mm, the Langenburg area 30 mm, the Goodeve area 37 mm, the Kelvington area 34 mm, the Foam Lake area 13 mm, the

Bulyea area 21 mm and the Semans area 11 mm. Cereal crops in many of the fields that received a lot of rain are now lodged.

Estimated East-central Hay Yields (tons/acre) - July 16, 2012		
	Dry land	Irrigated Land
Alfalfa	1.5	2.2
Brome/Alfalfa	1.7	2.2
Other Tame Hay	1.4	1.9
Wild Hay	1.4	N/A
Greenfeed	2.6	2.5

Cropland topsoil moisture is rated as 18 per cent surplus, 81 per cent adequate and one per cent short. Hay land and pasture topsoil moisture is rated as 25 per cent surplus, 74 per cent adequate and one per cent short. Crop District 5A is reporting 53 per cent of cropland acres and 47 per cent of hay land and pasture acres to have surplus moisture.

Livestock producers now have 52 per cent of the hay crop cut and 31 per cent baled or put into silage. Hay quality is rated as 20 per cent excellent, 71 per cent good and nine per cent fair. Current conditions are ideal for haying, although there are indications that some hay crops will not be harvested due to excess moisture.

Fungicides are being applied to most crops in the area and there are reports of some fields being sprayed for wheat midge. The majority of crop damage this week is due to flooding, wind, insects and disease.

Farmers are busy haying, controlling disease and insects and getting equipment ready for harvest.

West-Central Saskatchewan (Crop Districts 6B and 7)

Warm temperatures are quickly advancing crop development in the area, although there are reports that some crops are still significantly behind for this time of year. Varying amounts of rainfall were received this past week, with the Wilkie area reporting the greatest amount (102 mm). The Outlook area received 27 mm of rainfall, the Langham

area 66 mm, the Netherhill area 40 mm, the Kindersley area 16 mm, the Kerrobert area 65 mm and the Major area 22 mm.

Topsoil moisture conditions are rated as 29 per cent surplus and 71 per cent adequate on cropland. Hay land and pasture topsoil moisture conditions are rated as 23 per cent surplus and 77 per cent adequate. Crop District 7B is reporting 38 per cent of cropland has surplus of topsoil moisture.

Estimated West-central Hay Yields (tons/acre) - July 16, 2012		
	Dry land	Irrigated Land
Alfalfa	2.0	3.1
Brome/Alfalfa	1.8	3.5
Other Tame Hay	1.3	3.0
Wild Hay	1.4	N/A
Greenfeed	2.0	3.5

Haying operations progressed quickly with the warm weather, and livestock producers have 58 per cent of the hay crop cut and 30 per cent baled or put into silage. Hay quality is rated as 27 per cent excellent, 58 per cent good and 15 per cent fair. High humidity is making haying difficult. There are indications that hay yields will be less than last year.

Thunderstorms brought hail and heavy rain to the region, lodging some crops. Fungicides and insecticides are being applied to crops, and there are reports that most applications have been done by airplane. There are concerns about wheat midge and bertha armyworm populations in some parts of the area. The majority of crop damage this past week was due to flooding, hail, wind and diseases.

Producers are busy haying, controlling disease and hauling grain.

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

Heavy rain and strong winds have lodged many crops in the region. High temperatures have stressed canola crops and additional moisture and cooler weather would be welcome. The Vonda area received 82 mm of rain this past week, the Star City area 68 mm, the Nipawin area 40 mm, the Humboldt area 30 mm, the Melfort area 69 mm and the Garrick area 33 mm. Since April 1, the Kinistino area has received 416 mm of precipitation.

Estimated Northeast Hay Yields (tons/acre) - July 16, 2012		
	Dry land	Irrigated Land
Alfalfa	1.5	N/A
Brome/Alfalfa	1.4	N/A
Other Tame Hay	1.1	N/A
Wild Hay	1.1	N/A
Greenfeed	0.8	N/A

Topsoil moisture on cropland is rated as 39 per cent surplus and 61 per cent adequate. Topsoil moisture on hay land and pasture is reported as 24 per cent surplus and 76 per cent adequate. Crop districts 8A and 8B are reporting 48 per cent and 42 per cent of their respective cropland acres to have surplus moisture.

Livestock producers now have 72 per cent of the hay crop cut and 52 per cent baled or put into silage. Hay quality is rated as 25 per cent excellent and 75 per cent good. Most crop damage this week is attributed to flooding, wind and disease. Many crops are being sprayed with fungicides, and there are reports that some fields are being sprayed for wheat midge.

Producers are busy haying, controlling disease and hauling grain.

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

Rainfall across most of the region this past week has helped alleviate heat stress on the crops, although there are reports of heavy lodging in some areas due to heavy rain. The Hafford area received 65 mm of rainfall, the North Battleford area 44 mm, the

Neilburg area 25 mm, the St. Walburg area 18 mm and the Barthel area 3 mm. Some northern areas could use some additional moisture to help advance crops.

Estimated Northwest Hay Yields (tons/acre) - July 16, 2012		
	Dry land	Irrigated Land
Alfalfa	1.7	N/A
Brome/Alfalfa	1.7	N/A
Other Tame Hay	1.2	N/A
Wild Hay	1.1	N/A
Greenfeed	2.2	N/A

Topsoil moisture on cropland is rated as six per cent surplus, 87 per cent adequate and seven per cent short. Topsoil moisture on hay land and pasture is reported as four per cent surplus, 86 per cent adequate and 10 per cent short.

Haying is advancing quickly and livestock producers now have 48 per cent of the hay crop cut and 25 per cent baled or put into silage. Hay quality is rated as 35 per cent excellent and 65 per cent good. Wet conditions have delayed haying operations in some areas.

The majority of crop damage this week is attributed to flooding, wind and drought. Crops are being sprayed for leaf disease.

Producers are busy haying and controlling disease.

Crop Conditions – July 16, 2012

SK (provincial) Crop conditions - July 16, 2012							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	26	37	20	18	13	15	14
% good	61	58	62	63	66	64	66
% fair	11	5	16	18	19	17	18
% poor	1	0	2	1	2	3	2
% very poor	1	0	0	0	0	1	0
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	19	18	29	18	11	15	2
% good	55	62	56	61	62	69	88
% fair	20	18	13	18	22	15	10
% poor	5	2	2	3	4	1	0
% very poor	1	0	0	0	1	0	0

Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	27	37	18	18	13	10	7
% good	59	55	53	51	65	66	60
% fair	12	8	28	30	19	23	32
% poor	1	0	1	0	2	1	0
% very poor	1	0	0	1	1	0	1
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	18	3	12	10	7	5	0
% good	52	35	49	51	42	64	90
% fair	28	62	34	33	47	31	10
% poor	1	0	4	5	1	0	0
% very poor	1	0	1	1	3	0	0

Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	27	42	18	18	11	10	13
% good	63	54	64	68	73	76	70
% fair	10	4	17	13	16	13	17
% poor	0	0	1	1	0	1	0
% very poor	0	0	0	0	0	0	0
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	11	16	30	20	10	15	N/A
% good	73	59	58	66	77	70	N/A
% fair	15	22	11	12	9	15	N/A
% poor	1	3	1	2	4	0	N/A
% very poor	0	0	0	0	0	0	N/A


East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	23	28	19	31	10	17	19
% good	65	70	66	64	68	64	72
% fair	10	2	12	5	19	16	8
% poor	1	0	2	0	2	2	1
% very poor	1	0	1	0	1	1	0
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	19	39	35	35	19	47	N/A
% good	56	58	53	57	66	50	N/A
% fair	17	3	10	8	11	3	N/A
% poor	6	0	2	0	1	0	N/A
% very poor	2	0	0	0	3	0	N/A

West-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	30	31	21	16	11	15	7
% good	68	62	61	60	71	68	75
% fair	2	7	17	23	18	16	17
% poor	0	0	1	1	0	1	1
% very poor	0	0	0	0	0	0	0
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	19	9	23	13	14	0	10
% good	60	73	62	59	69	70	80
% fair	17	16	12	24	15	15	10
% poor	4	2	2	4	2	15	0
% very poor	0	0	1	0	0	0	0

Northeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	11	0	17	33	7	8	26
% good	69	100	57	46	62	53	45
% fair	17	0	21	19	27	28	15
% poor	3	0	5	1	4	10	13
% very poor	0	0	0	1	0	1	1
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	14	0	19	0	1	N/A	N/A
% good	43	50	50	70	55	N/A	N/A
% fair	29	35	17	18	27	N/A	N/A
% poor	11	10	11	7	17	N/A	N/A
% very poor	3	5	3	5	0	N/A	N/A

Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Flax
% excellent	0	0	27	N/A	32	29	0
% good	93	92	66	N/A	63	63	100
% fair	7	8	6	N/A	5	8	0
% poor	0	0	1	N/A	0	0	0
% very poor	0	0	0	N/A	0	0	0
	Canola	Mustard	Pea	Lentil	Canaryseed	Chickpea	Sunflower
% excellent	32	N/A	46	0	N/A	N/A	N/A
% good	58	N/A	50	100	N/A	N/A	N/A
% fair	10	N/A	4	0	N/A	N/A	N/A
% poor	0	N/A	0	0	N/A	N/A	N/A
% very poor	0	N/A	0	0	N/A	N/A	N/A

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period July 10 to 16, 2012

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	N/A	181	4A	19	Frontier	6	194.6	7B	347 A	Biggar	45	379
	2	Mount Pleasant	N/A	166		49	White Valley	N/A	217		347 B	Biggar	38	337
	3	Enniskillen	26	284		51	Reno	20.2	187.2		350 A	Mariposa	65.2	410.4
	33	Moose Creek	N/A	207		79	Arlington	10	215		350 B	Mariposa	47	287
	34	Browning	N/A	199		109	Carmichael	6	288		351	Progress	32	347
	61 A	Autler	N/A	256		110	Papot	3	244		352	Heart's Hill	22	366
	61 B	Autler	N/A	NIL		111	Maple Creek	11	296		377	Glenade	60	404
	62	Moose Mountain	22	286.3	4B	139	Gull Lake	8	298		378 A	Rosemount	45	373
	64	Brock	13	273		141	Big Stick	7	313		378 B	Rosemount	63	378
	65	Tecumseh	7	242		142	Enterprise	16	266		379	Reford	69	360
1B	91	Maryfield	26	318		169	Pittville	N/A	186		381	Grass Lake	N/A	316
	92	Walpole	N/A	249		231	Happyland	N/A	234		409 A	Buffalo	102	450
	95	Golden West	15	257	5A	183	Fertile Belt	15	419		409 B	Buffalo	N/A	234.6
	122	Marlin	20	370		186	Abemethy	16	323		410	Round Valley	37.5	269.2
	123	Silverwood	17	325		211	Churchbridge	30	359		411	Senlac	N/A	274
	124	Kingsley	15	240		213	Salcoats	19	340	BA	395	Porcupine	57	336
	151 A	Rocanville	26	476		214	Cana	N/A	N/A		397	Barner Valley	N/A	253
	151 B	Rocanville	10	417		216	Tullymet	N/A	230		428	Star City	68	374
	153	Wilowdale	12	349		217	Lupton	16	316		456	Arboret	27	331
	154	Elkapo	11	248		241	Calder	51	341		457	Connaught	38	393
	155	Wohaley	16.5	336		243	Wallace	43	307		458	Willow Creek	50	279
2A	66	Griffin	N/A	N/A		245 A	Garry	33	420		488	Moose Range	N/A	258
	67	Weyburn	N/A	177		245 B	Garry	29	423		487	Nipawin	40	317
	68	Brokenshell	7	210		245 C	Garry	36.5	339	BB	369	St. Peter	45	309
	69	Norton	N/A	N/A		246	Iuna Bon Accord	21	291		370	Humboldt	30	326
	97	Wellington	15	242		247	Kellors	28	306		371	Bayne	44	295
2B	127 A	Francis	15.5	277.5		248	Touchwood	16	223		372	Grant	82	337.5
	127 B	Francis	22	230.5	5B	271	Cole	26	405		402	Fish Creek	48	314
	129	Bratt's Lake	15.5	275.5		276	Foam Lake	11	288.5		429	Flett's Springs	69	341
	131	Baldon	18	325		277	Emerald	14	383		459	Kindino	80	416
	156	Indian Head	13.5	329		305	Invermay	31	326		460	Birch Hills	48	356.4
	160 A	Pense	23	284		307	Elros	20	367	SAE	461	Prince Albert	N/A	241.7
	160 B	Pense	22	281.5		308	Big Quill	N/A	240		488	Torch River	33	233
	161	Moose Jaw	19	321		333 A	Clayton	40	258.9		491	Buckland	N/A	250.9
	162	Caron	37	338.7		334	Preeceville	43	313	SAW	406	Mayfield	57	383
	191	Margus	28	286		336	Sarnan	31	335		435	Redberry	65	372
3ASE	38 A	Laurier	14	225.2		337	Lakeview	30	274		436	Douglas	48	260
	38 B	Laurier	14	199		338	Lakeside	19	213		463	Duck Lake	58	348.5
	39 A	The Gap	N/A	164		366	Kelvington	34	316		464	Leask	N/A	15
	39 B	The Gap	8	209	6A	189	Lumsden	10	272		467 A	Round Hill	44	235
3ASW	10	Happy Valley	7	219		190 A	Dufferin	6	269		467 B	Round Hill	23	261
	12	Poplar Valley	3	271		190 B	Dufferin	17	270		494	Camwood	21.5	243
	40	Bengough	9	181		190 C	Dufferin	4	220.5		497	Medstead	N/A	195
	42	Willow Bunch	15	287		190 D	Dufferin	N/A	305.6	9B	438	Battle River	N/A	283
	43	Old Post	5	248		219	Longlakeston	23	229		440	Hillsdale	29	256
	44	Waverley	N/A	245.5		220	McKellop	21	283		442	Manitou Lake	25.1	258.4
	70	Key West	N/A	160		221	Sarnia	10.4	274		498 B	Meda	N/A	38
	71	Excel	8	193		222	Craik	9	306		498 A	Parkdale	45	314
	73 A	Stonehenge	5	197.3		251	Big Arm	27	249		498 B	Parkdale	35	243
	73 B	Stonehenge	5	242		252	Arm River	10	252		490	Mervin	23	246
	74	Wood River	NIL	174.3		279	Mount Hope	11	248		501 A	Frenchman Butte	10	249
3AN	101	Termit	46	285		282	McCraney	24	309		501 B	Frenchman Butte	18	187
	102	Lake Johnston	7	232.7		309	Prairie Rose	6	245		501 C	Frenchman Butte	11	266
	103	Sutton	NIL	253		313	Lost River	33	273		561	Loon Lake	3	210
	104	Gravelbourg	10	227.5		339	Leroy	28	364.1		588 A	Meadow Lake	1	159
	132	Hillsborough	49	355.5		340	Wolverine	40	290		588 B	Meadow Lake	1	217
	134	Shamrock	8	225		341	Viscount	43	315		588 C	Meadow Lake	1	169
	164	Chaplin	8	227		343	Blucher	32	307.5		588 D	Meadow Lake	NIL	237.1
	193 A	Eyebrow	18	277		283	Rosedale	10	290.5		622	Beaver River	N/A	206
	193 B	Eyebrow	37	372.5	6B	284	Rusby	27	405					
	194	Entinist	N/A	303		285	Fertile Valley	21	392					
3BS	17	Vai Marie	13	421		286	Milden	N/A	324					
	18	Lone Tree	N/A	N/A		314	Dundum	19	311					
	45	Manikita	NIL	306		344	Coman Park	66	347					
	75 A	Pinto Creek	8	336		346	Perdise	36	401					
	75 B	Pinto Creek	NIL	241		376 A	Eagle Creek	62	350					
	76 A	Auvergne	3	293		376 B	Eagle Creek	46	458					
	76 B	Auvergne	N/A	282		403	Rotherham	50	341					
	77	Wise Creek	4	250		259	Slope Lake	N/A	316.1					
	78 A	Grassy Creek	8	255	7A	260 A	Newcombe	N/A	238					
	78 B	Grassy Creek	N/A	210		260 B	Newcombe	13	296					
	78 C	Grassy Creek	12	226		287	St. Andrews	27	372					
	105	Glenbair	3	263		288	Pleasant Valley	28	354					
	106	Whiska Creek	7	299		290	Kandensley	40	262					
	107	Lac Pelletier	3	259		292	Milton	25	316.5					
	108	Bone Creek	N/A	241		317 A	Marmot	50	320					
3BN	138	Webb	7	300		317 B	Marmot	39.5	351					
	166	Exchehor	15	318		318	Mountain View	21	367					
	167	Sark Landing	9.2	272.4		320 A	Oakdale	24	276					
	168	Riverside	5	221		320 B	Oakdale	16	235					
	226	Victory	N/A	337		321	Prairieville	13	286					
	228	Lacadena	10	250.5										
	257	Monet	8	252.7										



Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Weekly Rainfall

for the week ending July 16, 2012

Rainfall (mm)




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Saskatchewan
Ministry of
Agriculture

© 2012 Government of Saskatchewan



Projection: UTM Zone 13 Datum: NAD83



Data Source:


Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: July 18, 2012

Cumulative Rainfall

From: April 1, 2012

To: July 16, 2012



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Saskatchewan
Ministry of
Agriculture

0 25 50 100 150 200
Kilometers



© 2012 Government of Saskatchewan

Projection UTM Zone 13 Datum NAD83



Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: July 18, 2012

Cropland Topsoil Moisture Conditions

July 17, 2012

Moisture Conditions




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Saskatchewan
Ministry of
Agriculture

© 2012 Government of Saskatchewan



Projection: UTM Zone 13 Datum: NAD83




Data Source:

Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: July 18, 2012

Hay and Pasture Topsoil Moisture Conditions

July 17, 2012



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Saskatchewan
Ministry of
Agriculture

© 2012 Government of Saskatchewan

0 25 50 100 150 200
Kilometers

Projection: UTM Zone 13 Datum: NAD83



Data Source:

Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: July 18, 2012